Table 2

 Common Multiplication and Division Situations\*\*

|   | Unknown Product  | Group Size Unknown ("How many in each group?" Division)  | Number of Groups Unknown ("How many groups?" Division)   |
|---|--|--|--|
|   | 3 <b>•</b> 6 = ?   | $3 \bullet ? = 18$ , and $18 \div 3 = ?$   | $? \bullet 6 = 18$ , and $18 \div 6 = ?$   |
| Equal Groups                                    | There are 3 bags with 6 plums in each bag. How many plums are there in all?  | If 18 plums are shared equally into 3 bags, then how many plums will be in each bag?   | If 18 plums are to be packed 6 to a bag, then how many bags are needed?  |
|   | Measurement example. You need 3 lengths of string, each 6 inches long. How much string will you need altogether?                       | Measurement example. You have 18 inches of string, which you will cut into 3 equal pieces. How long will each piece of string be?                                | Measurement example. You have 18 inches of string, which you will cut into pieces that are 6 inches long. How many pieces of string will you have?                                   |
| Arrays <sup>4</sup><br>and<br>Area <sup>5</sup> | There are 3 rows of apples with 6 apples in each row. How many apples are there?   | If 18 apples are arranged into 3 equal rows, how many apples will be in each row?  | If 18 apples are arranged into equal rows of 6 apples, how many rows will there be?  |
|   | Area example. What is the area of a rectangle that is 3 centimeters by 6 centimeters?  | Area example. A rectangle has area 18 square centimeters. If one side is 3 centimeters long, how long is a side next to it?                                      | Area example. A rectangle has area 18 square centimeters. If one side is 6 centimeters long, how long is a side next to it?  |
| Compare   | A blue hat costs \$6. A red hat costs 3 times as much as the blue hat. How much does the red hat cost?                                 | A red hat costs \$18 and that is 3 times as much as a blue hat costs. How much does a blue hat cost?   | A red hat costs \$18 and a blue hat costs \$6. How many times as much does the red hat cost as the blue hat?   |
|   | Measurement example. A rubber band is 6 centimeters long. How long will the rubber band be when it is stretched to be 3 times as long? | Measurement example. A rubber band is stretched to be 18 centimeters long and that is 3 times as long as it was at first. How long was the rubber band at first? | Measurement example. A rubber band was 6 centimeters long at first. Now it is stretched to be 18 centimeters long. How many times as long is the rubber band now as it was at first? |
| General   | <i>a</i> • <i>b</i> = ?  | $a \bullet ? = p$ , and $p \div a = ?$   | $? \bullet b = p$ , and $p \div b = ?$   |

<sup>&</sup>lt;sup>4</sup>The language in the array examples shows the easiest form of array problems. A harder form is to use the terms rows and columns: The apples in the grocery window are in 3 rows and 6 columns. How many apples are in there? Both forms are valuable.

<sup>&</sup>lt;sup>5</sup>Area involves arrays of squares that have been pushed together so that there are no gaps or overlaps, so array problems include these especially important measurement situations.

<sup>\*\*</sup>The first examples in each cell are examples of discrete things. These are easier for students and should be given before the measurement examples.